
## Administering Influenza Vaccines: Intramuscular and Intranasal

## Intramuscular (IM) Injection

Inactivated Influenza Vaccines (aIIV3, HD-IIV3, IIV4, ccIIV4, RIV4)

- Use a needle long enough to reach deep into the muscle. For infants ≥6 months and children, use a 1" needle. For adolescents and adults, a 1-1 ½" needle should be used.¹
- 2. Choose the appropriate site. With your left hand<sup>2</sup>, spread the skin taut between the thumb and forefinger, isolating the muscle.
- 3. With your right hand<sup>2</sup>, insert the needle at a 90° angle to the skin with a quick thrust.
- 4. Push down on the plunger and inject the entire contents of the syringe. There is no need to aspirate.



- 5. Withdraw the needle and apply light pressure to the injection site for several seconds with a gauze pad.
- 6. If there is any bleeding, cover the injection site with a bandage.
- 7. Put used syringe in a sharps container.


## Intranasal (IN/NAS) Administration

Live Attenuated Influenza Vaccine (LAIV4)

- 1. FluMist® (LAIV4) is for intranasal administration only. Do not inject FluMist®.
- 2. Remove the rubber tip protector. Do not remove the dose-divider clip at the other end of the sprayer.
- 3. With the patient in an upright position, place the tip just inside the nostril to ensure LAIV is delivered into the nose. The patient should breathe normally.



- 4. With a single motion, depress the plunger as rapidly as possible until the dose-divider clip prevents you from going further. Remove from nostril.
- 5. Pinch and remove the dose-divider clip from the plunger.



- 6. Place the tip just inside the other nostril, and with a single motion, depress plunger as rapidly as possible to deliver the remaining vaccine.
- 7. Put applicator in a sharps container.

For more information regarding flu and flu vaccination, refer to <a href="www.michigan.gov/flu">www.cdc.gov/vaccines</a>, or <a href="www.cdc.gov/mmwr">www.cdc.gov/vaccines</a>, or <a href="www.cdc.gov/mmwr">www.cdc.gov/mmwr</a>.

<sup>&</sup>lt;sup>1</sup> Use professional judgment when determining needle size and injection site. For more information, see General Best Practice Guidelines for Immunization at <a href="https://www.cdc.gov/vaccines/hcp/acip-recs/general-recs/index.html">www.cdc.gov/vaccines/hcp/acip-recs/general-recs/index.html</a>.

<sup>&</sup>lt;sup>2</sup> Use the opposite hand if you are left-handed.